

EIGENVALUES OF $\mathcal{PT}$ -SYMMETRIC OSCILLATORS WITH POLYNOMIAL POTENTIALS

KWANG C. SHIN

ABSTRACT. We study the eigenvalue problem $-u''(z) - [(iz)^m + P_{m-1}(iz)]u(z) = \lambda u(z)$ with the boundary conditions that $u(z)$ decays to zero as z tends to infinity along the rays $\arg z = -\frac{\pi}{2} \pm \frac{2\pi}{m+2}$, where $P_{m-1}(z) = a_1 z^{m-1} + a_2 z^{m-2} + \cdots + a_{m-1} z$ is a polynomial and integers $m \geq 3$. We provide an asymptotic expansion of the eigenvalues λ_n as $n \rightarrow +\infty$, and prove that for each *real* polynomial P_{m-1} , the eigenvalues are all real and positive, with only finitely many exceptions.

2000 Mathematics subject classification: 34L40, 34L20

1. INTRODUCTION

For integers $m \geq 3$ fixed, we are considering the “non-standard” non-self-adjoint eigenvalue problems

$$(1) \quad Hu(z, \lambda) := \left[-\frac{d^2}{dz^2} - (iz)^m - P_{m-1}(iz) \right] u(z, \lambda) = \lambda u(z, \lambda), \quad \text{for some } \lambda \in \mathbb{C},$$

with the boundary condition that

$$(2) \quad u(z, \lambda) \rightarrow 0 \text{ exponentially, as } z \rightarrow \infty \text{ along the two rays } \arg(z) = -\frac{\pi}{2} \pm \frac{2\pi}{m+2},$$

where P_{m-1} is a polynomial of degree at most $m-1$ of the form

$$(3) \quad P_{m-1}(z) = a_1 z^{m-1} + a_2 z^{m-2} + \cdots + a_{m-1} z, \quad a_j \in \mathbb{C} \text{ for } 1 \leq j \leq m-1.$$

We let

$$a := (a_1, a_2, \dots, a_{m-1}) \in \mathbb{C}^{m-1}$$

be the coefficient vector of $P_{m-1}(z)$. We are mainly interested in the case when P_{m-1} is real, that is, when $a \in \mathbb{R}^{m-1}$. However, some interesting facts in this paper hold also for $a \in \mathbb{C}^{m-1}$. So except for Theorem 4 below, we will use $a \in \mathbb{C}^{m-1}$.

If a nonconstant function u satisfies (1) with some $\lambda \in \mathbb{C}$ and the boundary condition (2), then we call λ an *eigenvalue* of H and u an *eigenfunction of H associated with the eigenvalue λ* . Also, the *geometric multiplicity of an eigenvalue λ* is the number of linearly independent eigenfunctions associated with the eigenvalue λ . The operator H in (1) with potential $V(z) = -(iz)^m - P_{m-1}(iz)$ is called *$\mathcal{PT}$ -symmetric* if $\overline{V(-\bar{z})} = V(z)$, $z \in \mathbb{C}$. Note that $V(z) = -(iz)^m - P_{m-1}(iz)$ is a $\mathcal{PT}$ -symmetric potential if and only if $a \in \mathbb{R}^{m-1}$.

Before we state our main theorems, we first introduce some known facts by Sibuya [18] about the eigenvalues λ of H .

Theorem 1. *The eigenvalues λ_n of H have the following properties.*

- (I) *The set of all eigenvalues is a discrete set in $\mathbb{C}$.*
- (II) *The geometric multiplicity of every eigenvalue is one.*
- (III) *Infinitely many eigenvalues, accumulating at infinity, exist.*
- (IV) *The eigenvalues have the following asymptotic expansion*

$$(4) \quad \lambda_n = \left(\frac{\Gamma\left(\frac{3}{2} + \frac{1}{m}\right) \sqrt{\pi} \left(n - \frac{1}{2}\right)}{\sin\left(\frac{\pi}{m}\right) \Gamma\left(1 + \frac{1}{m}\right)} \right)^{\frac{2m}{m+2}} [1 + o(1)] \quad \text{as } n \text{ tends to infinity, } n \in \mathbb{N},$$

where the error term $o(1)$ could be complex-valued.

This paper is organized as follows. In Section 2, we will introduce work of Hille [12] and Sibuya [18], regarding properties of solutions of (1). We then improve on the asymptotics of a certain function in [18]. In Section 3, we introduce an entire function $C(a, \lambda)$ whose zeros are the eigenvalues of H , due to Sibuya [18]. In Section 4, we then provide asymptotics of $C(a, \lambda)$ as $\lambda \rightarrow \infty$ in the complex plane, improving the asymptotics of $C(a, \lambda)$ in [18]. In Section 5, we will improve the asymptotic expansion (4) of the eigenvalues. In particular, we will prove the following. Throughout this paper, we use that $\lfloor x \rfloor$ is the largest integer that is less than or equal to $x \in \mathbb{R}$.

Theorem 2. *Let $a \in \mathbb{C}^{m-1}$ be fixed. Then there exist $e_\ell(a) \in \mathbb{C}$, $1 \leq \ell \leq \frac{m}{2} + 1$ such that the eigenvalues λ_n of H have the asymptotic expansion*

$$(5) \quad \lambda_n \underset{n \rightarrow +\infty}{=} \lambda_{0,n} + \sum_{\ell=1}^{\lfloor \frac{m}{2} + 1 \rfloor} e_\ell(a) \lambda_{0,n}^{1 - \frac{\ell}{m}} + o\left(\lambda_{0,n}^{\frac{1}{2} - \frac{1}{m}}\right),$$

where

$$\lambda_{0,n} = \left(\frac{\left(n + \frac{1}{2}\right) \pi}{K_m \sin\left(\frac{2\pi}{m}\right)} \right)^{\frac{2m}{m+2}} \quad \text{with} \quad K_m = \int_0^\infty \left(\sqrt{1+t^m} - \sqrt{t^m} \right) dt > 0.$$

One can compute K_m directly (or see equation (2.22) in [9] with the identity $\Gamma(s)\Gamma(1-s) = \pi \csc(\pi s)$) and obtains

$$K_m = \frac{\sqrt{\pi} \Gamma\left(1 + \frac{1}{m}\right)}{2 \cos\left(\frac{\pi}{m}\right) \Gamma\left(\frac{3}{2} + \frac{1}{m}\right)}.$$

In the last section, we prove the following theorem, regarding monotonicity of $|\lambda_n|$.

Theorem 3. *For each $a \in \mathbb{C}^{m-1}$ there exists $M > 0$ such that $|\lambda_n| < |\lambda_{n+1}|$ if $n \geq M$.*

This is a consequence of (5).

Finally, when H is $\mathcal{PT}$ -symmetric (i.e., $a \in \mathbb{R}^{m-1}$), $u(z, \lambda)$ is an eigenfunction associated with an eigenvalue λ if and only if $\overline{u(-\bar{z}, \lambda)}$ is an eigenfunction associated with the eigenvalue

$\bar{\lambda}$. Thus, the eigenvalues either appear in complex conjugate pairs, or else are real. So Theorem 3 implies the following.

Theorem 4. *Suppose that $a \in \mathbb{R}^{m-1}$. Then the eigenvalues λ of H are all real and positive, with only finitely many exceptions.*

For the rest of the Introduction, we will mention a brief history of problem (1).

In recent years, these $\mathcal{PT}$ -symmetric operators have gathered considerable attention, because ample numerical and asymptotic studies suggest that many of such operators have real eigenvalues only even though they are not self-adjoint. In particular, the differential operators H with some polynomial potential V and with the boundary condition (2) have been considered by Bessis and Zinn-Justin (not in print), Bender and Boettcher [2] and many other physicists [3, 4, 5, 9, 13, 14, 15, 17, 19].

Around 1992 Bessis and Zinn-Justin (not in print) conjectured that when $V(z) = iz^3 + \beta z^2$, $\beta \in \mathbb{R}$, the eigenvalues are all real and positive, and in 1998, Bender and Boettcher [2] conjectured that when $V(z) = (iz)^m + \beta z^2$, $\beta \in \mathbb{R}$, the eigenvalues are all real and positive. Many numerical, asymptotic and analytic studies support these conjectures (see, e. g., [3, 4, 5, 9, 13, 14, 15, 17, 19] and references therein and below).

The first rigorous proof of reality and positivity of the eigenvalues of some non-self-adjoint H in (1) was given by Dorey, Dunning and Tateo [8] in 2001. They proved that the eigenvalues of H with the potential $V(z) = -(iz)^{2m} - \alpha(iz)^{m-1} + \frac{\ell(\ell+1)}{z^2}$, $m, \alpha, \ell \in \mathbb{R}$, are all real if $m > 1$ and $\alpha < m + 1 + |2\ell + 1|$, and positive if $m > 1$ and $\alpha < m + 1 - |2\ell + 1|$.

Then in 2002 the present author [16] extended the polynomial potential results of Dorey, Dunning and Tateo to more general polynomial cases, by adapting the method in [8]. Namely, when $V(z) = -(iz)^m - P_{m-1}(iz)$, the eigenvalues are all real and positive, provided that for some $1 \leq j \leq \frac{m}{2}$ the coefficients of the real polynomial P_{m-1} satisfy $(j - k)a_k \geq 0$ for all $1 \leq k \leq m - 1$.

However, there are some $\mathcal{PT}$ -symmetric polynomial potentials that produce non-real eigenvalues. Delabaere and Pham [6], and Delabaere and Trinh [7] studied the potential $iz^3 + \gamma iz$ and showed that a pair of non-real eigenvalues develops for large negative γ . Moreover, Handy [10], and Handy, Khan, Wang and Tymczak [11] showed that the same potential admits a pair of non-real eigenvalues for small negative values of $\gamma \approx -3.0$. Also, Bender, Berry, Meisinger, Savage and Simsek [1] considered the problem with the potential $V(z) = z^4 + iAz$, $A \in \mathbb{R}$, under decaying boundary conditions at both ends of the real axis, and their numerical study showed that more and more non-real eigenvalues develop as $|A| \rightarrow \infty$. So without any restrictions on the coefficients a_k , Theorem 4 is the most general result one can expect about reality of eigenvalues.




FIGURE 1. The Stokes sectors for $m = 3$. The dashed rays represent $\arg z = \pm\frac{\pi}{5}, \pm\frac{3\pi}{5}, \pi$.

Also, the method used to prove Theorem 4 in this paper is new. The method used in [8, 16] is useful in proving reality of *all* eigenvalues, but I think that some critical arguments in proving reality of eigenvalues in [8, 16] cannot be applied to the cases when some non-real eigenvalues exist. The asymptotic expansion (5) itself is interesting, and also (5) implies Theorem 3. Note that (4) is not enough to conclude Theorem 3. Finally, Theorem 3 and $\mathcal{PT}$ -symmetry of H explained right before Theorem 4 above imply the partial reality of the eigenvalues in Theorem 4.

2. PROPERTIES OF THE SOLUTIONS

In this section, we introduce work of Hille [12] and Sibuya [18] about properties of the solutions of (1).

First, we scale equation (1) because many facts that we need later are stated for the scaled equation. Let u be a solution of (1) and let $v(z, \lambda) = u(-iz, \lambda)$. Then v solves

$$(6) \quad -v''(z, \lambda) + [z^m + P_{m-1}(z) + \lambda]v(z, \lambda) = 0,$$

where $m \geq 3$ and P_{m-1} is a polynomial (possibly, $P_{m-1} \equiv 0$) of the form (3).

Since we scaled the argument of u , we must rotate the boundary conditions. We state them in a more general context by using the following definition.

Definition. The Stokes sectors S_k of the equation (6) are

$$S_k = \left\{ z \in \mathbb{C} : \left| \arg(z) - \frac{2k\pi}{m+2} \right| < \frac{\pi}{m+2} \right\} \quad \text{for } k \in \mathbb{Z}.$$

See Figure 1. It is known from Hille [12, §7.4] that every nonconstant solution of (6) either decays to zero or blows up exponentially, in each Stokes sector S_k . That is, one has the following result.

Lemma 5 ([12, §7.4]).

- (i) For each $k \in \mathbb{Z}$, every solution v of (6) (with no boundary conditions imposed) is asymptotic to

$$(7) \quad (\text{const.}) z^{-\frac{m}{4}} \exp \left[\pm \int^z [\xi^m + P_{m-1}(\xi) + \lambda]^{\frac{1}{2}} d\xi \right]$$

as $z \rightarrow \infty$ in every closed subsector of S_k .

- (ii) If a nonconstant solution v of (6) decays in S_k , it must blow up in $S_{k-1} \cup S_{k+1}$. However, when v blows up in S_k , v need not be decaying in S_{k-1} or in S_{k+1} .

Lemma 5 (i) implies that if v decays along one ray in S_k , then it decays along all rays in S_k . Also, if v blows up along one ray in S_k , then it blows up along all rays in S_k . Thus, since the rotation $z \mapsto iz$ maps the two rays in (2) onto the center rays of S_{-1} and S_1 ,

the boundary conditions on u in (1) mean that v decays in $S_{-1} \cup S_1$.

Next we will introduce Sibuya's results, but first we define a sequence of complex numbers b_j in terms of the a_k and λ , as follows. For $\lambda \in \mathbb{C}$ fixed, we expand

$$(8) \quad \begin{aligned} & (1 + a_1 z^{-1} + a_2 z^{-2} + \cdots + a_{m-1} z^{1-m} + \lambda z^{-m})^{1/2} \\ &= 1 + \sum_{k=1}^{\infty} \binom{\frac{1}{2}}{k} (a_1 z^{-1} + a_2 z^{-2} + \cdots + a_{m-1} z^{1-m} + \lambda z^{-m})^k \\ &= 1 + \sum_{j=1}^{\infty} \frac{b_j(a, \lambda)}{z^j}, \quad \text{for large } |z|. \end{aligned}$$

Note that $b_1, b_2, \dots, b_{m-1}$ do not depend on λ , so we write $b_j(a) = b_j(a, \lambda)$ for $j = 1, 2, \dots, m-1$. So the above expansion without the λz^{-m} term still gives b_j for $1 \leq j \leq m-1$. We further define $r_m = -\frac{m}{4}$ if m is odd, and $r_m = -\frac{m}{4} - b_{\frac{m}{2}+1}(a)$ if m is even.

The following theorem is a special case of Theorems 6.1, 7.2, 19.1 and 20.1 of Sibuya [18] that is the main ingredient of the proofs of the main results in this paper.

Theorem 6. Equation (6), with $a \in \mathbb{C}^{m-1}$, admits a solution $f(z, a, \lambda)$ with the following properties.

- (i) $f(z, a, \lambda)$ is an entire function of z, a and λ .
(ii) $f(z, a, \lambda)$ and $f'(z, a, \lambda) = \frac{\partial}{\partial z} f(z, a, \lambda)$ admit the following asymptotic expansions.
Let $\varepsilon > 0$. Then

$$\begin{aligned} f(z, a, \lambda) &= z^{r_m} (1 + O(z^{-1/2})) \exp [-F(z, a, \lambda)], \\ f'(z, a, \lambda) &= -z^{r_m + \frac{m}{2}} (1 + O(z^{-1/2})) \exp [-F(z, a, \lambda)], \end{aligned}$$

as z tends to infinity in the sector $|\arg z| \leq \frac{3\pi}{m+2} - \varepsilon$, uniformly on each compact set of (a, λ) -values. Here

$$F(z, a, \lambda) = \frac{2}{m+2} z^{\frac{m}{2}+1} + \sum_{1 \leq j < \frac{m}{2}+1} \frac{2}{m+2-2j} b_j(a) z^{\frac{1}{2}(m+2-2j)}.$$

(iii) Properties (i) and (ii) uniquely determine the solution $f(z, a, \lambda)$ of (6).

(iv) For each fixed $a \in \mathbb{C}^{m-1}$ and $\delta > 0$, f and f' also admit the asymptotic expansions,

$$(9) \quad f(0, a, \lambda) = [1 + o(1)] \lambda^{-1/4} \exp [L(a, \lambda)],$$

$$(10) \quad f'(0, a, \lambda) = -[1 + o(1)] \lambda^{1/4} \exp [L(a, \lambda)],$$

as $\lambda \rightarrow \infty$ in the sector $|\arg(\lambda)| \leq \pi - \delta$, where

$$L(a, \lambda) = \begin{cases} \int_0^{+\infty} \left(\sqrt{t^m + P_{m-1}(t) + \lambda} - t^{\frac{m}{2}} - \sum_{j=1}^{\frac{m+1}{2}} b_j(a) t^{\frac{m}{2}-j} \right) dt & \text{if } m \text{ is odd,} \\ \int_0^{+\infty} \left(\sqrt{t^m + P_{m-1}(t) + \lambda} - t^{\frac{m}{2}} - \sum_{j=1}^{\frac{m}{2}} b_j(a) t^{\frac{m}{2}-j} - \frac{b_{\frac{m}{2}+1}(a)}{t+1} \right) dt & \text{if } m \text{ is even.} \end{cases}$$

(v) The entire functions $\lambda \mapsto f(0, a, \lambda)$ and $\lambda \mapsto f'(0, a, \lambda)$ have orders $\frac{1}{2} + \frac{1}{m}$.

Proof. In Sibuya's book [18], see Theorem 6.1 for a proof of (i) and (ii); Theorem 7.2 for a proof of (iii); and Theorem 19.1 for a proof of (iv). Moreover, (v) is a consequence of (iv) along with Theorem 20.1. Note that properties (i), (ii) and (iv) are summarized on pages 112–113 of Sibuya [18]. $\square$

Using this theorem, Sibuya [18, Theorem 19.1] also showed the following corollary that will be useful later on.

Corollary 7. *Let $a \in \mathbb{C}^{m-1}$ be fixed. Then $L(a, \lambda) = K_m \lambda^{\frac{1}{2} + \frac{1}{m}} (1 + o(1))$ as λ tends to infinity in the sector $|\arg \lambda| \leq \pi - \delta$, and hence*

$$(11) \quad \operatorname{Re} (L(a, \lambda)) = K_m \cos \left(\frac{m+2}{2m} \arg(\lambda) \right) |\lambda|^{\frac{1}{2} + \frac{1}{m}} (1 + o(1))$$

as $\lambda \rightarrow \infty$ in the sector $|\arg(\lambda)| \leq \pi - \delta$.

In particular, $\operatorname{Re} (L(a, \lambda)) \rightarrow +\infty$ as $\lambda \rightarrow \infty$ in any closed subsector of the sector $|\arg(\lambda)| < \frac{m\pi}{m+2}$. In addition, $\operatorname{Re} (L(a, \lambda)) \rightarrow -\infty$ as $\lambda \rightarrow \infty$ in any closed subsector of the sectors $\frac{m\pi}{m+2} < |\arg(\lambda)| < \pi - \delta$.

Proof. This asymptotic expansion will be clear from Lemma 8 below, or alternatively, see [18, Theorem 19.1] for a proof. $\square$

Based on the above Corollary, Sibuya [18, Theorem 29.1] also proved the following asymptotic expansion of the eigenvalues.

$$(12) \quad \lambda_n = \omega^m \left(\frac{(-2n+1)\pi}{2K_m \sin \left(\frac{2\pi}{m} \right)} \right)^{\frac{2m}{m+2}} [1 + o(1)], \quad \text{as } n \rightarrow \infty,$$

where

$$\omega = \exp \left[\frac{2\pi i}{m+2} \right].$$

Notice that in this paper we consider the boundary conditions of the scaled equation (6) where v decays in $S_{-1} \cup S_1$, while Sibuya studies equation (6) with boundary conditions such that v decays in $S_0 \cup S_2$. The factor ω^m in our formula (12) is due to this scaling of the problem.

Remark. Throughout this paper, we will deal with numbers like $(\omega^\nu \lambda)^s$ for some $s \in \mathbb{R}$, and $\nu \in \mathbb{C}$. As usual, we will use

$$\omega^\nu = \exp \left[\nu \frac{2\pi i}{m+2} \right]$$

and if $\arg(\lambda)$ is specified, then

$$\arg((\omega^\nu \lambda)^s) = s [\arg(\omega^\nu) + \arg(\lambda)] = s \left[\operatorname{Re}(\nu) \frac{2\pi}{m+2} + \arg(\lambda) \right], \quad s \in \mathbb{R}.$$

If $s \notin \mathbb{Z}$ then the branch of λ^s is chosen to be the negative real axis.

Next, we provide an improved asymptotic expansion of L . We will use this new asymptotic expansion of L to improve the asymptotic expansion (12) of the eigenvalues.

Lemma 8. *Let $m \geq 3$ and $a \in \mathbb{C}^{m-1}$ be fixed. Then there exist constants $K_{m,j}(a) \in \mathbb{C}$, $0 \leq j \leq \frac{m}{2} + 1$, such that*

$$L(a, \lambda) = \begin{cases} \sum_{j=0}^{\frac{m+1}{2}} K_{m,j}(a) \lambda^{\frac{1}{2} + \frac{1-j}{m}} + O\left(|\lambda|^{-\frac{1}{2m}}\right) & \text{if } m \text{ is odd,} \\ \sum_{j=0}^{\frac{m}{2}+1} K_{m,j}(a) \lambda^{\frac{1}{2} + \frac{1-j}{m}} - \frac{b_{\frac{m}{2}+1}(a)}{m} \ln(\lambda) + O\left(|\lambda|^{-\frac{1}{m}}\right) & \text{if } m \text{ is even,} \end{cases}$$

as $\lambda \rightarrow \infty$ in the sector $|\arg(\lambda)| \leq \pi - \delta$.

Proof. The function $L(a, \lambda)$ is defined as an integral over $0 \leq t < +\infty$ in Theorem 6. We will rotate the contour of integration using Cauchy's integral formula. In doing so, we need to justify that the integrand in the definition of $L(a, \lambda)$ is analytic in some domain in the complex plane.

Let $0 < \delta < \frac{\pi}{m+2}$ be a fixed number. Suppose that $0 \leq \arg(\lambda) \leq \pi - \delta$. Then if $0 \leq \arg(t) \leq \frac{1}{m} \arg(\lambda)$, there exists $M_0 > 0$ such that

$$-\pi < -\frac{\delta}{2} \leq \arg(t^m + P_{m-1}(t)) \leq \arg(\lambda) + \frac{\delta}{2} \leq \pi - \frac{\delta}{2},$$

provided that $|t| \geq M_0$. Since $t^m + P_{m-1}(t)$ lies in a large disk centered at the origin for $|t| \leq M_0$, we see that for all λ with $|\lambda|$ large, we have that $-\frac{\delta}{2} < \arg(t^m + P_{m-1}(t) + \lambda) < \pi - \frac{\delta}{2}$ and $|t^m + P_{m-1}(t) + \lambda| > 0$ for all t in the sector $0 \leq \arg(t) \leq \frac{1}{m} \arg(\lambda)$, and hence $\sqrt{t^m + P_{m-1}(t) + \lambda}$ is analytic in the sector $0 \leq \arg(t) \leq \frac{1}{m} \arg(\lambda)$ if λ lies outside a large disk and in the sector $0 \leq \arg(\lambda) \leq \pi - \delta$.

Let

$$Q(t, a, \lambda) = \begin{cases} \sqrt{t^m + P_{m-1}(t) + \lambda} - t^{\frac{m}{2}} - \sum_{j=1}^{\frac{m+1}{2}} b_j(a) t^{\frac{m}{2}-j} & \text{if } m \text{ is odd,} \\ \sqrt{t^m + P_{m-1}(t) + \lambda} - t^{\frac{m}{2}} - \sum_{j=1}^{\frac{m}{2}} b_j(a) t^{\frac{m}{2}-j} - \frac{b_{\frac{m}{2}+1}(a)}{t+1} & \text{if } m \text{ is even.} \end{cases}$$

Then, since $|Q(t, a, \lambda)| = O(|t|^{-\frac{m}{2}})$ as t tends to infinity in the sector $0 \leq \arg(t) \leq \frac{1}{m} \arg(\lambda)$, we have by Cauchy's integral formula, upon substituting $t = \lambda^{\frac{1}{m}} \tau$ for all λ with $|\lambda|$ large enough,

$$(13) \quad L(a, \lambda) = \int_0^{+\infty} Q(t, a, \lambda) dt = \lambda^{\frac{1}{m}} \int_0^{+\infty} Q(\lambda^{\frac{1}{m}} \tau, a, \lambda) d\tau,$$

where

$$Q(\lambda^{\frac{1}{m}} \tau, a, \lambda) = \begin{cases} \lambda^{\frac{1}{2}} \left(\sqrt{\tau^m + 1 + \frac{P_{m-1}(\lambda^{\frac{1}{m}} \tau)}{\lambda}} - \tau^{\frac{m}{2}} - \sum_{j=1}^{\frac{m+1}{2}} b_j(a) \frac{\tau^{\frac{m}{2}-j}}{\lambda^{\frac{j}{m}}} \right) & \text{if } m \text{ is odd,} \\ \lambda^{\frac{1}{2}} \left(\sqrt{\tau^m + 1 + \frac{P_{m-1}(\lambda^{\frac{1}{m}} \tau)}{\lambda}} - \tau^{\frac{m}{2}} - \sum_{j=1}^{\frac{m}{2}} b_j(a) \frac{\tau^{\frac{m}{2}-j}}{\lambda^{\frac{j}{m}}} - \frac{\lambda^{-\frac{1}{2}} b_{\frac{m}{2}+1}(a)}{\lambda^{\frac{1}{m}} \tau + 1} \right) & \text{if } m \text{ is even.} \end{cases}$$

Similarly, (13) holds for $-\pi + \delta \leq \arg(\lambda) \leq 0$.

Next, we examine the following square root in $Q(\lambda^{\frac{1}{m}} \tau, a, \lambda)$:

$$\begin{aligned} \sqrt{\tau^m + 1 + \frac{P_{m-1}(\lambda^{\frac{1}{m}} \tau)}{\lambda}} &= \sqrt{\tau^m + 1} \sqrt{1 + \frac{P_{m-1}(\lambda^{\frac{1}{m}} \tau)}{\lambda(\tau^m + 1)}} \\ &= \sqrt{\tau^m + 1} \left(1 + \sum_{k=1}^{\infty} \binom{\frac{1}{2}}{k} \left(\frac{P_{m-1}(\lambda^{\frac{1}{m}} \tau)}{\lambda(\tau^m + 1)} \right)^k \right) \\ &\stackrel{\text{let}}{=} \sqrt{\tau^m + 1} + \sum_{j=1}^{\infty} \frac{g_j(\tau)}{\lambda^{\frac{j}{m}}}, \end{aligned}$$

where $g_j(\tau)$ are functions such that $g_j(\tau)$ are all integrable on $[0, R]$ for any $R > 0$. Moreover, by the definition of b_j in (8), we see that for $1 \leq j \leq m-1$,

$$g_j(\tau) = \sum_{k=1}^j \frac{b_{j,k}(a) \tau^{mk-j}}{(\tau^m + 1)^{k-\frac{1}{2}}} \quad \text{for some constants } b_{j,k}(a) \text{ such that } \sum_{k=1}^j b_{j,k}(a) = b_j(a).$$

Thus,

$$\begin{aligned} g_j(\tau) - b_j(a) \tau^{\frac{m}{2}-j} &= \sum_{k=1}^j b_{j,k}(a) \left(\frac{\tau^{mk-j}}{(\tau^m + 1)^{k-\frac{1}{2}}} - \tau^{\frac{m}{2}-j} \right) \\ &\stackrel{\tau \rightarrow \infty}{=} \sum_{k=1}^j b_{j,k}(a) \tau^{\frac{m}{2}-j} O\left(\frac{1}{\tau^m}\right) \\ &\stackrel{\tau \rightarrow \infty}{=} O\left(\frac{1}{\tau^{\frac{m}{2}+j}}\right) \quad \text{for all } 1 \leq j \leq \frac{m+1}{2}. \end{aligned}$$

So $\int_0^\infty |g_j(\tau) - b_j(a)\tau^{\frac{m}{2}-j}| d\tau < +\infty$ for all $1 \leq j \leq \frac{m+1}{2}$. Next, when m is even and $j = \frac{m}{2} + 1$, we write

$$\begin{aligned} & \int_0^\infty \left(g_{\frac{m}{2}+1}(\tau) - \frac{b_{\frac{m}{2}+1}(a)}{\tau + \lambda^{-\frac{1}{m}}} \right) d\tau \\ &= \int_0^\infty \left(g_{\frac{m}{2}+1}(\tau) - \frac{b_{\frac{m}{2}+1}(a)}{\tau + 1} \right) d\tau + b_{\frac{m}{2}+1}(a) \int_0^\infty \left(\frac{1}{\tau + 1} - \frac{1}{\tau + \lambda^{-\frac{1}{m}}} \right) d\tau \\ &\stackrel{\text{let}}{=} K_{m, \frac{m}{2}+1}(a) - \frac{b_{\frac{m}{2}+1}(a)}{m} \ln(\lambda), \end{aligned}$$

where we take $\text{Im}(\ln(\lambda)) = \arg(\lambda) \in (-\pi, \pi)$.

Thus, we have that

$$L(a, \lambda) = \begin{cases} \sum_{j=0}^{\frac{m+1}{2}} K_{m,j}(a) \lambda^{\frac{1}{2} + \frac{1-j}{m}} + O\left(|\lambda|^{-\frac{1}{2m}}\right) & \text{if } m \text{ is odd,} \\ \sum_{j=0}^{\frac{m}{2}+1} K_{m,j}(a) \lambda^{\frac{1}{2} + \frac{1-j}{m}} - \frac{b_{\frac{m}{2}+1}(a)}{m} \ln(\lambda) + O\left(|\lambda|^{-\frac{1}{m}}\right) & \text{if } m \text{ is even,} \end{cases}$$

as $\lambda \rightarrow \infty$ in the sector $|\arg(\lambda)| \leq \pi - \delta$, where

$$\begin{aligned} (14) \quad & K_{m,0}(a) = K_m = \int_0^\infty \left(\sqrt{1+t^m} - \sqrt{t^m} \right) dt > 0 \quad \text{for all } m \geq 3, \\ & K_{m,j}(a) = \int_0^\infty \left(g_j(t) - b_j(a)t^{\frac{m}{2}-j} \right) dt \quad \text{for all } 1 \leq j \leq \frac{m+1}{2}, \\ & K_{m, \frac{m}{2}+1}(a) = \int_0^\infty \left(g_{\frac{m}{2}+1}(t) - \frac{b_{\frac{m}{2}+1}(a)}{t+1} \right) dt \quad \text{when } m \text{ is even.} \end{aligned}$$

This completes the proof. $\square$

3. EIGENVALUES ARE ZEROS OF AN ENTIRE FUNCTION

In this section, we will prove that the eigenvalues are zeros of an entire function.

First, we let

$$G^k(a) := (\omega^{-k}a_1, \omega^{-2k}a_2, \dots, \omega^{-(m-1)k}a_{m-1}) \quad \text{for } k \in \mathbb{Z}.$$

Then recall that the function $f(z, a, \lambda)$ in Theorem 6 solves (6) and decays to zero exponentially as $z \rightarrow \infty$ in S_0 , and blows up in $S_{-1} \cup S_1$. Next, one can check that the function

$$f_k(z, a, \lambda) := f(\omega^{-k}z, G^k(a), \omega^{-mk}\lambda),$$

which is obtained by scaling $f(z, G^k(a), \omega^{-mk}\lambda)$ in the z -variable, also solves (6). It is clear that $f_0(z, a, \lambda) = f(z, a, \lambda)$. Also, $f_k(z, a, \lambda)$ decays in S_k and blows up in $S_{k-1} \cup S_{k+1}$ since $f(z, G^k(a), \omega^{-mk}\lambda)$ decays in S_0 . Since no nonconstant solution decays in two consecutive Stokes sectors (see Lemma 5 (ii)), f_k and f_{k+1} are linearly independent and hence any solution of (6) can be expressed as a linear combination of these two. Especially, there exist some coefficients $C(a, \lambda)$ and $\tilde{C}(a, \lambda)$ such that

$$(15) \quad f_{-1}(z, a, \lambda) = C(a, \lambda)f_0(z, a, \lambda) + \tilde{C}(a, \lambda)f_1(z, a, \lambda).$$

We then see that

$$(16) \quad C(a, \lambda) = \frac{W_{-1,1}(a, \lambda)}{W_{0,1}(a, \lambda)} \quad \text{and} \quad \tilde{C}(a, \lambda) = -\frac{W_{-1,0}(a, \lambda)}{W_{0,1}(a, \lambda)},$$

where $W_{j,k} = f_j f'_k - f'_j f_k$ is the Wronskian of f_j and f_k . Since both f_j and f_k are solutions of the same linear equation (6), we know that the Wronskians are constant functions of z . Also, since f_k and f_{k+1} are linearly independent, $W_{k,k+1} \neq 0$ for all $k \in \mathbb{Z}$. Moreover, we have the following lemma that is useful later on.

Lemma 9. *Suppose $k, j \in \mathbb{Z}$. Then*

$$(17) \quad W_{k+1,j+1}(a, \lambda) = \omega^{-1} W_{k,j}(G(a), \omega^2 \lambda),$$

and $W_{0,1}(a, \lambda) = 2\omega^{\mu(a)}$, where

$$\mu(a) = \begin{cases} \frac{m}{4} & \text{if } m \text{ is odd,} \\ \frac{m}{4} - b_{\frac{m}{2}+1}(a) & \text{if } m \text{ is even.} \end{cases}$$

Moreover,

$$\tilde{C}(a, \lambda) = -\frac{W_{-1,0}(a, \lambda)}{W_{0,1}(a, \lambda)} = -\omega \frac{W_{0,1}(G^{-1}(a), \omega^{-2} \lambda)}{W_{0,1}(a, \lambda)} = -\omega^{1+2\nu(a)},$$

where

$$(18) \quad \nu(a) = \begin{cases} 0 & \text{if } m \text{ is odd,} \\ b_{\frac{m}{2}+1}(a) & \text{if } m \text{ is even.} \end{cases}$$

Proof. See Sibuya [18, pages 116-118] for proof. Here, we mention that by (8), we have $b_{\frac{m}{2}+1}(G^{-1}(a)) = -b_{\frac{m}{2}+1}(a)$ and hence $\nu(G^{-1}(a)) = -\nu(a)$. $\square$

Now we can identify the eigenvalues of H as the zeros of the entire function $\lambda \mapsto C(a, \lambda)$.

Theorem 10. *For each fixed $a \in \mathbb{C}^{m-1}$, the function $\lambda \mapsto C(a, \lambda)$ is entire. Moreover, λ is an eigenvalue of H if and only if $C(a, \lambda) = 0$.*

Proof. Since $W_{0,1}(a, \lambda) \neq 0$ and since $W_{-1,1}(a, \lambda)$ is a Wronskian of two entire functions, it is clear from (16) that $C(a, \lambda)$ is an entire function of λ for each fixed $a \in \mathbb{C}^{m-1}$.

Next, suppose that λ is an eigenvalue of H with a corresponding eigenfunction u , then the scaled eigenfunction $v(z, \lambda) = u(-iz, \lambda)$ solves (6) and decays in $S_{-1} \cup S_1$. Hence, v is a (nonzero) constant multiple of f_1 since both decays in S_1 . Similarly, v is also a constant multiple of f_{-1} . Thus, f_{-1} is a constant multiple of f_1 , implying $C(a, \lambda) = 0$.

Conversely, if $C(a, \lambda) = 0$, then f_{-1} is a constant multiple of f_1 , and hence f_1 also decays in S_{-1} . Thus, f_1 decays in $S_{-1} \cup S_1$ and is a scaled eigenfunction with the eigenvalue λ . $\square$

Moreover, the following is an easy consequence of (15): For each $k \in \mathbb{Z}$ we have

$$(19) \quad W_{-1,k}(a, \lambda) = C(a, \lambda) W_{0,k}(a, \lambda) + \tilde{C}(a) W_{1,k}(a, \lambda),$$

where we use $\tilde{C}(a)$ for $\tilde{C}(a, \lambda)$ since it is independent of λ .

4. ASYMPTOTIC EXPANSIONS OF $C(a, \lambda)$

In this section, we provide asymptotic expansions of the entire function $C(a, \lambda)$ as $\lambda \rightarrow \infty$ along all possible rays to infinity in the complex plane.

First, we provide an asymptotic expansion of the Wronskian of f_0 and f_j in preparation for providing an asymptotic expansion of $C(a, \lambda)$.

Lemma 11. *Suppose that $1 \leq j \leq \frac{m}{2} + 1$. Then for each $a \in \mathbb{C}^{m-1}$,*

$$(20) \quad W_{0,j}(a, \lambda) = [2i\omega^{-\frac{j}{2}} + o(1)] \exp [L(G^j(a), \omega^{2j-m-2}\lambda) + L(a, \lambda)],$$

as $\lambda \rightarrow \infty$ in the sector

$$(21) \quad -\pi + \delta \leq \pi - \frac{4j\pi}{m+2} + \delta \leq \arg(\lambda) \leq \pi - \delta.$$

Proof. We fix $1 \leq j \leq \frac{m}{2} + 1$. Then,

$$\begin{aligned} W_{0,j}(a, \lambda) &= f_0(z, a, \lambda)f_j'(z, a, \lambda) - f_0'(z, a, \lambda)f_j(z, a, \lambda) \\ &= \omega^{-j}f(0, a, \lambda)f'(0, G^j(a), \omega^{2j-m-2}\lambda) - f'(0, a, \lambda)f(0, G^j(a), \omega^{2j-m-2}\lambda) \\ &\underset{\lambda \rightarrow \infty}{=} - \left[\omega^{-j}\omega^{\frac{2j-m-2}{4}} - \omega^{-\frac{2j-m-2}{4}} + o(1) \right] \exp [L(G^j(a), \omega^{2j-m-2}\lambda) + L(a, \lambda)] \\ &\underset{\lambda \rightarrow \infty}{=} [2i\omega^{-\frac{j}{2}} + o(1)] \exp [L(G^j(a), \omega^{2j-m-2}\lambda) + L(a, \lambda)], \end{aligned}$$

where we used (9) and (10) with

$$|\arg(\lambda)| \leq \pi - \delta \quad \text{and} \quad |\arg(\omega^{2j-m-2}\lambda)| \leq \pi - \delta,$$

which is, (21). Here we also used $j \leq \frac{m}{2} + 1$. $\square$

Next, we provide an asymptotic expansion of $W_{-1,1}(a, \lambda)$ as $\lambda \rightarrow \infty$ along the rays near the negative real axis. Notice from (16) that $W_{-1,1}(a, \lambda) = W_{0,1}(a, \lambda)C(a, \lambda)$. Also, $W_{0,1}(a, \lambda)$ is a nonzero constant function of λ . So from these one gets an asymptotic expansion of $C(a, \lambda)$.

Theorem 12. *For each fixed $a \in \mathbb{C}^{m-1}$ and $0 < \delta < \frac{\pi}{m+2}$,*

$$(22) \quad W_{-1,1}(a, \lambda) = [2i + o(1)] \exp [L(G^{-1}(a), \omega^{-2}\lambda) + L(G(a), \omega^{-m}\lambda)],$$

as $\lambda \rightarrow \infty$ along the rays in the sector

$$(23) \quad \pi - \frac{4\pi}{m+2} + \delta \leq \arg(\lambda) \leq \pi + \frac{4\pi}{m+2} - \delta.$$

Moreover, there exists a constant $M_1 > 0$ such that $W_{-1,1}(a, \lambda) \neq 0$ for all λ in the sector

$$(23) \text{ if } |\lambda| \geq M_1.$$

Proof. This is an easy consequence of Lemma 11 and equation (17).

The last assertion of the theorem is a consequence of the asymptotic expansion (22). $\square$

The asymptotic expansion of $C(a, \lambda)$ in a sector near the positive real axis is obtained in the following theorem.

Theorem 13. Suppose that $m \geq 4$. Then for each fixed $a \in \mathbb{C}^{m-1}$ and $0 < \delta < \frac{\pi}{m+2}$,

$$C(a, \lambda) = [\omega^{\frac{1}{2}} + o(1)] \exp [L(G^{-1}(a), \omega^{-2}\lambda) - L(a, \lambda)] \\ + [\omega^{\frac{1}{2}+2\nu(a)} + o(1)] \exp [L(G(a), \omega^2\lambda) - L(a, \lambda)],$$

as $\lambda \rightarrow \infty$ in the sector

$$(24) \quad \pi - \frac{4\lfloor \frac{m}{2} \rfloor \pi}{m+2} + \delta \leq \arg(\lambda) \leq \pi - \frac{4\pi}{m+2} - \delta.$$

Proof. Suppose $2 \leq k \leq \frac{m}{2}$. Then from (17), (19) and Lemma 11,

$$\begin{aligned} C(a, \lambda) &= \frac{W_{-1,k}(a, \lambda)}{W_{0,k}(a, \lambda)} - \tilde{C}(a) \frac{W_{1,k}(a, \lambda)}{W_{0,k}(a, \lambda)} \\ &= \frac{\omega W_{0,k+1}(G^{-1}(a), \omega^{-2}\lambda)}{W_{0,k}(a, \lambda)} - \tilde{C}(a) \frac{\omega^{-1} W_{0,k-1}(G(a), \omega^2\lambda)}{W_{0,k}(a, \lambda)} \\ &= [\omega^{\frac{1}{2}} + o(1)] \frac{\exp [L(G^k(a), \omega^{2k-m-2}\lambda) + L(G^{-1}(a), \omega^{-2}\lambda)]}{\exp [L(G^k(a), \omega^{2k-m-2}\lambda) + L(a, \lambda)]} \\ &\quad - [\omega^{-\frac{1}{2}} + o(1)] \tilde{C}(a) \frac{\exp [L(G^k(a), \omega^{2k-m-2}\lambda) + L(G(a), \omega^2\lambda)]}{\exp [L(G^k(a), \omega^{2k-m-2}\lambda) + L(a, \lambda)]} \\ &= [\omega^{\frac{1}{2}} + o(1)] \exp [L(G^{-1}(a), \omega^{-2}\lambda) - L(a, \lambda)] \\ &\quad + [\omega^{-\frac{1}{2}} + o(1)] \omega^{1+2\nu(a)} \exp [L(G(a), \omega^2\lambda) - L(a, \lambda)], \end{aligned}$$

as $\lambda \rightarrow \infty$ such that

$$\begin{aligned} -\pi &< \pi - \frac{4(k+1)\pi}{m+2} + \delta \leq \arg(\omega^{-2}\lambda) \leq \pi - \delta, \\ \pi - \frac{4k\pi}{m+2} + \delta &\leq \arg(\lambda) \leq \pi - \delta, \\ \pi - \frac{4(k-1)\pi}{m+2} + \delta &\leq \arg(\omega^2\lambda) \leq \pi - \delta, \end{aligned}$$

that is,

$$\pi - \frac{4k\pi}{m+2} + \delta \leq \arg(\lambda) \leq \pi - \frac{4\pi}{m+2} - \delta,$$

provided that $2 \leq k \leq \frac{m}{2}$. So in order to complete the proof, we choose

$$k = \begin{cases} \frac{m-1}{2} & \text{if } m \text{ is odd and } m \geq 5, \\ \frac{m}{2} & \text{if } m \text{ is even.} \end{cases}$$

□

The sectors (23) and (24) do not cover the entire complex plane near infinity. The next theorem covers a sector in the upper half plane, connecting the sectors (23) and (24) in the upper half plane.

Theorem 14. Suppose that $a \in \mathbb{C}^{m-1}$ and $0 < \delta < \frac{\pi}{m+2}$. If $m \geq 4$ then

$$(25) \quad C(a, \lambda) = \left[\omega^{\frac{1}{2}} + o(1) \right] \exp [L(G^{-1}(a), \omega^{-2}\lambda) - L(a, \lambda)] \\ - [i\omega^{1+\mu(a)+4\nu(a)} + o(1)] \exp [-L(G^2(a), \omega^{2-m}\lambda) - L(a, \lambda)],$$

as $\lambda \rightarrow \infty$ in the sector

$$(26) \quad \pi - \frac{8\pi}{m+2} + \delta \leq \arg(\lambda) \leq \pi - \delta.$$

If $m = 3$ then

$$\begin{aligned} C(a, \lambda) = & [-\omega^{-2} + o(1)] \exp [L(G^4(a), \omega^{-2}\lambda) - L(a, \lambda)] \\ & - [i\omega^{\frac{7}{4}} + o(1)] \exp [-L(G^2(a), \omega^{-1}\lambda) - L(a, \lambda)], \end{aligned}$$

as $\lambda \rightarrow \infty$ in the sector

$$(27) \quad -\frac{\pi}{5} + \delta \leq \arg(\lambda) \leq \pi - \delta.$$

Moreover, if $m \geq 6$ then there exists a constant $M_2 > 0$ such that $C(a, \lambda) \neq 0$ for all λ in the sector (26) if $|\lambda| \geq M_2$.

Proof. Suppose that $m \geq 4$. Then from (17), (19) and Lemma 11,

$$\begin{aligned} C(a, \lambda) &= \frac{W_{-1,2}(a, \lambda)}{W_{0,2}(a, \lambda)} - \tilde{C}(a) \frac{W_{1,2}(a, \lambda)}{W_{0,2}(a, \lambda)} \\ &= \frac{\omega W_{0,3}(G^{-1}(a), \omega^{-2}\lambda)}{W_{0,2}(a, \lambda)} - \tilde{C}(a) \frac{\omega^{-1} W_{0,1}(G(a), \omega^2\lambda)}{W_{0,2}(a, \lambda)} \\ &= \frac{\omega [2i\omega^{-\frac{3}{2}} + o(1)] \exp [L(G^2(a), \omega^{2-m}\lambda) + L(G^{-1}(a), \omega^{-2}\lambda)]}{[2i\omega^{-\frac{2}{2}} + o(1)] \exp [L(G^2(a), \omega^{2-m}\lambda) + L(a, \lambda)]} \\ &\quad - \tilde{C}(a) \frac{\omega^{-1} W_{0,1}(G(a), \omega^2\lambda)}{[2i\omega^{-1} + o(1)] \exp [L(G^2(a), \omega^{2-m}\lambda) + L(a, \lambda)]} \\ &= [\omega^{\frac{1}{2}} + o(1)] \exp [L(G^{-1}(a), \omega^{-2}\lambda) - L(a, \lambda)] \\ &\quad - \frac{-\omega^{1+2\nu(a)} 2\omega^{\mu(G(a))}}{[2i + o(1)] \exp [L(G^2(a), \omega^{2-m}\lambda) + L(a, \lambda)]} \end{aligned}$$

as $\lambda \rightarrow \infty$ such that

$$-\pi + \delta \leq \pi - \frac{12\pi}{m+2} + \delta \leq \arg(\omega^{-2}\lambda) \leq \pi - \delta \quad \text{and} \quad \pi - \frac{8\pi}{m+2} + \delta \leq \arg(\lambda) \leq \pi - \delta,$$

that is,

$$\pi - \frac{8\pi}{m+2} + \delta \leq \arg(\lambda) \leq \pi - \delta.$$

Next, we use $2\nu(a) + \mu(G(a)) = \mu(a) + 4\nu(a)$ to get (25).

Suppose $m = 3$. Then $\omega^5 = 1$. Also, $W_{-3,0}(a, \lambda) = W_{2,0}(a, \lambda)$ since $f_{-3}(z, a, \lambda) = f_2(z, a, \lambda)$. Thus, we have that

$$\begin{aligned}
C(a, \lambda) &= \frac{W_{-1,2}(a, \lambda)}{W_{0,2}(a, \lambda)} - \tilde{C}(a) \frac{W_{1,2}(a, \lambda)}{W_{0,2}(a, \lambda)} \\
&= \frac{\omega^{-2}W_{-3,0}(G^2(a), \omega^4\lambda)}{W_{0,2}(a, \lambda)} - \tilde{C}(a) \frac{\omega^{-1}W_{0,1}(G(a), \omega^2\lambda)}{W_{0,2}(a, \lambda)} \\
&= -\frac{\omega^{-2}W_{0,2}(G^2(a), \omega^4\lambda)}{W_{0,2}(a, \lambda)} - \tilde{C}(a) \frac{\omega^{-1}W_{0,1}(G(a), \omega^2\lambda)}{W_{0,2}(a, \lambda)} \\
&= -\frac{\omega^{-2}W_{0,2}(G^2(a), \omega^{-1}\lambda)}{W_{0,2}(a, \lambda)} - \tilde{C}(a) \frac{\omega^{-1}W_{0,1}(G(a), \omega^2\lambda)}{W_{0,2}(a, \lambda)} \\
&= -\frac{\omega^{-2}[2i\omega^{-\frac{2}{2}} + o(1)] \exp[L(G^4(a), \omega^{-2}\lambda) + L(G^2(a), \omega^{-1}\lambda)]}{[2i\omega^{-\frac{2}{2}} + o(1)] \exp[L(G^2(a), \omega^{-1}\lambda) + L(a, \lambda)]} \\
&\quad - \tilde{C}(a) \frac{\omega^{-1}W_{0,1}(G(a), \omega^2\lambda)}{[2i\omega^{-1} + o(1)] \exp[L(G^2(a), \omega^{-1}\lambda) + L(a, \lambda)]} \\
&= [-\omega^{-2} + o(1)] \exp[L(G^4(a), \omega^{-2}\lambda) - L(a, \lambda)] \\
&\quad - \frac{-\omega^{1+2\nu(a)} 2\omega^{\mu(G(a))}}{[2i + o(1)] \exp[L(G^2(a), \omega^{2-m}\lambda) + L(a, \lambda)]},
\end{aligned}$$

as $\lambda \rightarrow \infty$ such that

$$-\pi + \delta \leq \pi - \frac{8\pi}{5} + \delta \leq \arg(\omega^{-1}\lambda) \leq \pi - \delta \quad \text{and} \quad \pi - \frac{8\pi}{5} + \delta \leq \arg(\lambda) \leq \pi - \delta,$$

that is,

$$\pi - \frac{6\pi}{5} + \delta \leq \arg(\lambda) \leq \pi - \delta.$$

In order to show the last assertion, we suppose that $C(a, \lambda) = 0$ for some λ in (26) with large $|\lambda|$. Then from the asymptotic expansion (25), we have

$$(28) \quad \exp[L(G^{-1}(a), \omega^{-2}\lambda) + L(G^2(a), \omega^{2-m}\lambda)] = [i\omega^{-\frac{1}{2}+\mu(a)+4\nu(a)} + o(1)].$$

By Corollary 7,

$$\begin{aligned}
\operatorname{Re} (L(G^{-1}(a), \omega^{-2}\lambda)) &= K_m \cos \left(\frac{m+2}{2m} \arg(\omega^{-2}\lambda) \right) |\lambda|^{\frac{1}{2}+\frac{1}{m}} (1 + o(1)) \\
&= K_m \cos \left(-\frac{2\pi}{m} + \frac{m+2}{2m} \arg(\lambda) \right) |\lambda|^{\frac{1}{2}+\frac{1}{m}} (1 + o(1)), \\
\operatorname{Re} (L(G^2(a), \omega^{2-m}\lambda)) &= K_m \cos \left(\frac{m+2}{2m} \arg(\omega^{2-m}\lambda) \right) |\lambda|^{\frac{1}{2}+\frac{1}{m}} (1 + o(1)) \\
&= -K_m \cos \left(\frac{2\pi}{m} + \frac{m+2}{2m} \arg(\lambda) \right) |\lambda|^{\frac{1}{2}+\frac{1}{m}} (1 + o(1)).
\end{aligned}$$

Note that if $m \geq 6$, then $0 < \delta \leq \arg(\lambda) \leq \pi - \delta$ in (26). Since

$$\begin{aligned} & \cos\left(-\frac{2\pi}{m} + \frac{m+2}{2m} \arg(\lambda)\right) - \cos\left(\frac{2\pi}{m} + \frac{m+2}{2m} \arg(\lambda)\right) \\ &= 2 \sin\left(\frac{2\pi}{m}\right) \sin\left(\frac{m+2}{2m} \arg(\lambda)\right) > 0, \end{aligned}$$

we see that

$$\operatorname{Re} \left(L(G^{-1}(a), \omega^{-2}\lambda) + L(G^2(a), \omega^{2-m}\lambda) \right) \rightarrow +\infty,$$

as $\lambda \rightarrow \infty$ in (26), and hence the left hand side of (28) blows up. Thus, $C(a, \lambda)$ cannot have infinitely many zeros in (26). This completes the proof. $\square$

The next theorem covers a sector in the lower half plane, connecting sectors (23) and (24).

Theorem 15. *Suppose that $a \in \mathbb{C}^{m-1}$ and $0 < \delta < \frac{\pi}{m+2}$. If $m \geq 4$ then*

$$\begin{aligned} C(a, \lambda) &= [-i\omega^{1+\mu(a)} + o(1)] \exp [-L(a, \omega^{-m-2}\lambda) - L(G^{-2}(a), \omega^{-4}\lambda)] \\ &\quad + [\omega^{\frac{1}{2}+2\nu(a)} + o(1)] \exp [L(G(a), \omega^{-m}\lambda) - L(a, \omega^{-m-2}\lambda)], \end{aligned}$$

as $\lambda \rightarrow \infty$ in the sector

$$(29) \quad \pi + \delta \leq \arg(\lambda) \leq \pi + \frac{8\pi}{m+2} - \delta.$$

If $m = 3$ then

$$\begin{aligned} C(a, \lambda) &= [-i\omega^{\frac{7}{4}} + o(1)] \exp [-L(a, \omega^{-5}\lambda) - L(G^{-2}(a), \omega^{-4}\lambda)] \\ &\quad + [\omega^3 + o(1)] \exp [L(G(a), \omega^{-3}\lambda) - L(a, \omega^{-5}\lambda)], \end{aligned}$$

as $\lambda \rightarrow \infty$ in the sector

$$(30) \quad \pi + \delta \leq \arg(\lambda) \leq 2\pi + \frac{\pi}{5} - \delta.$$

Moreover, if $m \geq 6$ then there exists a constant $M_3 > 0$ such that $C(a, \lambda) \neq 0$ for all λ in the sector (29) if $|\lambda| \geq M_3$.

Proof. Suppose that $m \geq 4$. Then from (17), (19) and Lemma 11,

$$\begin{aligned}
C(a, \lambda) &= \frac{W_{-1,-2}(a, \lambda)}{W_{0,-2}(a, \lambda)} - \tilde{C}(a) \frac{W_{1,-2}(a, \lambda)}{W_{0,-2}(a, \lambda)} \\
&= \frac{W_{0,1}(G^{-2}(a), \omega^{-4}\lambda)}{W_{0,2}(G^{-2}(a), \omega^{-4}\lambda)} + \tilde{C}(a) \frac{\omega^{-1}W_{0,-3}(G(a), \omega^2\lambda)}{\omega^2 W_{0,2}(G^{-2}(a), \omega^{-4}\lambda)} \\
&= \frac{W_{0,1}(G^{-2}(a), \omega^{-4}\lambda)}{[2i\omega^{-\frac{2}{2}} + o(1)] \exp[L(a, \omega^{-m-2}\lambda) + L(G^{-2}(a), \omega^{-4}\lambda)]} \\
&\quad - \tilde{C}(a) \frac{[2i\omega^{-\frac{3}{2}} + o(1)] \exp[L(G(a), \omega^{-m}\lambda) + L(G^{-2}(a), \omega^{-4}\lambda)]}{[2i\omega^{-\frac{2}{2}} + o(1)] \exp[L(a, \omega^{-m-2}\lambda) + L(G^{-2}(a), \omega^{-4}\lambda)]} \\
&= \frac{2\omega^{\mu(G^{-2}(a))}}{[2i\omega^{-1} + o(1)] \exp[L(a, \omega^{-m-2}\lambda) + L(G^{-2}(a), \omega^{-4}\lambda)]} \\
&\quad + [\omega^{-\frac{1}{2}} + o(1)] \omega^{1+2\nu(a)} \frac{\exp[L(G(a), \omega^{-m}\lambda) + L(G^{-2}(a), \omega^{-4}\lambda)]}{\exp[L(a, \omega^{-m-2}\lambda) + L(G^{-2}(a), \omega^{-4}\lambda)]},
\end{aligned}$$

as $\lambda \rightarrow \infty$ such that

$$(31) \quad \pi - \frac{12\pi}{m+2} + \delta \leq \arg(\omega^{-4}\lambda) \leq \pi - \delta \quad \text{and} \quad \pi - \frac{8\pi}{m+2} + \delta \leq \arg(\omega^{-4}\lambda) \leq \pi - \delta,$$

that is,

$$(32) \quad \pi + \delta \leq \arg(\lambda) \leq \pi + \frac{8\pi}{m+2} - \delta,$$

which is (29).

Suppose that $m = 3$. Then,

$$\begin{aligned}
C(a, \lambda) &= \frac{W_{-1,-2}(a, \lambda)}{W_{0,-2}(a, \lambda)} - \tilde{C}(a) \frac{W_{1,-2}(a, \lambda)}{W_{0,-2}(a, \lambda)} \\
&= \frac{W_{0,1}(G^{-2}(a), \omega^{-4}\lambda)}{W_{0,2}(G^{-2}(a), \omega^{-4}\lambda)} + \tilde{C}(a) \frac{\omega^{-1}W_{0,-3}(G(a), \omega^2\lambda)}{\omega^2 W_{0,2}(G^{-2}(a), \omega^{-4}\lambda)} \\
&= \frac{W_{0,1}(G^{-2}(a), \omega^{-4}\lambda)}{W_{0,2}(G^{-2}(a), \omega^{-4}\lambda)} - \omega^2 \tilde{C}(a) \frac{W_{0,2}(G(a), \omega^{-3}\lambda)}{W_{0,2}(G^{-2}(a), \omega^{-4}\lambda)} \\
&= \frac{W_{0,1}(G^{-2}(a), \omega^{-4}\lambda)}{[2i\omega^{-\frac{2}{2}} + o(1)] \exp[L(a, \omega^{-5}\lambda) + L(G^{-2}(a), \omega^{-4}\lambda)]} \\
&\quad - \omega^2 \tilde{C}(a) \frac{[2i\omega^{-\frac{2}{2}} + o(1)] \exp[L(G^3(a), \omega^{-4}\lambda) + L(G(a), \omega^{-3}\lambda)]}{[2i\omega^{-\frac{2}{2}} + o(1)] \exp[L(a, \omega^{-5}\lambda) + L(G^{-2}(a), \omega^{-4}\lambda)]} \\
&= \frac{2\omega^{\mu(G^{-2}(a))}}{[2i\omega^{-1} + o(1)] \exp[L(a, \omega^{-5}\lambda) + L(G^{-2}(a), \omega^{-4}\lambda)]} \\
&\quad + [\omega^2 + o(1)] \omega^{1+2\nu(a)} \frac{\exp[L(G^{-2}(a), \omega^{-4}\lambda) + L(G(a), \omega^{-3}\lambda)]}{\exp[L(a, \omega^{-5}\lambda) + L(G^{-2}(a), \omega^{-4}\lambda)]},
\end{aligned}$$

as $\lambda \rightarrow \infty$ such that

$$\pi - \frac{8\pi}{5} + \delta \leq \arg(\omega^{-3}\lambda) \leq \pi - \delta \quad \text{and} \quad \pi - \frac{8\pi}{5} + \delta \leq \arg(\omega^{-4}\lambda) \leq \pi - \delta,$$

that is,

$$\pi + \delta \leq \arg(\lambda) \leq \pi + \frac{6\pi}{5} - \delta.$$

Finally, the proof of the last assertion of this theorem follows as in the proof of Theorem 14. $\square$

From the asymptotic expansions in the previous four theorems, one obtains the order of the entire function $\lambda \mapsto C(a, \lambda)$. The order of an entire function g is defined by

$$\limsup_{r \rightarrow \infty} \frac{\log \log M(r, g)}{\log r},$$

where $M(r, g) = \max\{|g(re^{i\theta})| : 0 \leq \theta \leq 2\pi\}$ for $r > 0$. If for some positive real numbers σ, c_1, c_2 , we have $\exp[c_1 r^\sigma] \leq M(r, g) \leq \exp[c_2 r^\sigma]$ for all large r , then the order of g is σ .

Corollary 16. *The entire function $\lambda \mapsto C(a, \lambda)$ is of order $\frac{1}{2} + \frac{1}{m}$.*

Proof. The sectors in (21), (23), (26), (29), cover a neighborhood of infinity in the complex plane. So the nonconstant entire function $|C(a, \lambda)|$ is bounded above by $\exp\left[c_1 |\lambda|^{\frac{1}{2} + \frac{1}{m}}\right]$ for some constant $c_1 > 0$. Also, along the ray $\arg(\lambda) = \pi$, one can see from (11) and (22) that $|C(a, \lambda)|$ is bounded below by $\exp\left[c_2 |\lambda|^{\frac{1}{2} + \frac{1}{m}}\right]$ for some constant $c_2 > 0$. Hence, the order of $C(a, \cdot)$ is $\frac{1}{2} + \frac{1}{m}$. $\square$

Remark. Since the eigenvalues are the zeros of the entire function $\lambda \mapsto C(a, \lambda)$ of order $\frac{1}{2} + \frac{1}{m} \in (0, 1)$, there are infinitely many discrete eigenvalues as was already mentioned in Theorem 1.

5. ASYMPTOTIC EXPANSION OF THE EIGENVALUES: PROOF OF THEOREM 2

In this section, we prove Theorem 2 by using the asymptotic expansions of $C(a, \lambda)$ and $L(a, \lambda)$.

Proof of Theorem 2. Recall that by Theorem 10, λ is an eigenvalue of H if and only if $C(a, \lambda) = 0$.

For $m \geq 4$ and $a \in \mathbb{C}^{m-1}$ fixed, suppose that $C(a, \lambda) = 0$ for some λ with $|\lambda|$ large. Then from the asymptotic expansion of $C(a, \lambda)$ in Theorem 13 we have

$$[1 + o(1)] \exp [L(G(a), \omega^2 \lambda) - L(G^{-1}(a), \omega^{-2} \lambda)] = -\omega^{-2\nu(a)},$$

and absorbing $[1 + o(1)]$ into the exponential function then yields

$$\exp [L(G(a), \omega^2 \lambda) - L(G^{-1}(a), \omega^{-2} \lambda) + o(1)] = -\omega^{-2\nu(a)}.$$

Thus, from Lemma 8 if m is odd, we infer

$$\begin{aligned}
& \ln(-\omega^{-2\nu(a)}) \\
&= L(G(a), \omega^2 \lambda) - L(G^{-1}(a), \omega^{-2} \lambda) + o(1) \\
(33) \quad &= \sum_{j=0}^{\lfloor \frac{m}{2} + 1 \rfloor} \left[K_{m,j}(G(a))(\omega^2 \lambda)^{\frac{1}{2} + \frac{1-j}{m}} - K_{m,j}(G^{-1}(a))(\omega^{-2} \lambda)^{\frac{1}{2} + \frac{1-j}{m}} \right] + o(1) \\
&= 2iK_{m,0} \sin\left(\frac{2\pi}{m}\right) \lambda^{\frac{1}{2} + \frac{1}{m}} + \sum_{j=1}^{\lfloor \frac{m}{2} + 1 \rfloor} c_{m,j}(a) \lambda^{\frac{1}{2} + \frac{1-j}{m}} + o(1),
\end{aligned}$$

where

$$(34) \quad c_{m,j}(a) = K_{m,j}(G(a))(\omega^2)^{\frac{1}{2} + \frac{1-j}{m}} - K_{m,j}(G^{-1}(a))(\omega^{-2})^{\frac{1}{2} + \frac{1-j}{m}}, \quad 1 \leq j \leq \frac{m+1}{2}.$$

Similarly, if m is even, then from Lemma 8 we have (33) with $c_{m,j}(a)$ in (34) for $1 \leq j \leq \frac{m}{2}$, and

$$c_{m, \frac{m}{2} + 1}(a) = K_{m, \frac{m}{2} + 1}(G(a)) - K_{m, \frac{m}{2} + 1}(G^{-1}(a)) + \frac{b_{\frac{m}{2} + 1}(a)}{m} \frac{8\pi i}{m+2},$$

where we used $b_{\frac{m}{2} + 1}(G^{-1}(a)) = -b_{\frac{m}{2} + 1}(a) = b_{\frac{m}{2} + 1}(G(a))$.

Note that there exist constants $M > 0$ and $\varepsilon > 0$ such that the function

$$(35) \quad \lambda \mapsto L(G(a), \omega^2 \lambda) - L(G^{-1}(a), \omega^{-2} \lambda) + o(1)$$

is continuous in the region $|\lambda| \geq M$ and $|\arg(\lambda)| \leq \varepsilon$. From (33) we then see that the function (35) maps the region $|\lambda| \geq M$ and $|\arg(\lambda)| \leq \varepsilon$ onto a region that contains the entire positive imaginary axis near infinity.

Thus, from (33) we get that for every sufficiently large $n \in \mathbb{N}$ there exists λ_n such that

$$2iK_{m,0} \sin\left(\frac{2\pi}{m}\right) \lambda_n^{\frac{1}{2} + \frac{1}{m}} + \sum_{j=1}^{\lfloor \frac{m}{2} + 1 \rfloor} c_{m,j}(a) \lambda_n^{\frac{1}{2} + \frac{1-j}{m}} + o(1) = \left(2n + 1 - \frac{4\nu(a)}{m+2}\right) \pi i.$$

Thus,

$$\lambda_n^{\frac{1}{2} + \frac{1}{m}} + \sum_{j=1}^{\lfloor \frac{m}{2} + 1 \rfloor} \frac{c_{m,j}(a)}{2iK_{m,0} \sin\left(\frac{2\pi}{m}\right)} \lambda_n^{\frac{1}{2} + \frac{1-j}{m}} + o(1) = \frac{\left(2n + 1 - \frac{4\nu(a)}{m+2}\right) \pi}{2K_{m,0} \sin\left(\frac{2\pi}{m}\right)}.$$

Let

$$(36) \quad d_{m,j}(a) = \begin{cases} \frac{c_{m,j}(a)}{2iK_{m,0} \sin\left(\frac{2\pi}{m}\right)} & \text{if } 1 \leq j \leq \frac{m}{2} + 1, \\ \frac{c_{m,j}(a) + \frac{4\nu(a)}{m+2} \pi i}{2iK_{m,0} \sin\left(\frac{2\pi}{m}\right)} & \text{if } m \text{ is even and } j = \frac{m}{2} + 1. \end{cases}$$

Then

$$(37) \quad \lambda_n^{\frac{1}{2} + \frac{1}{m}} + \sum_{j=1}^{\lfloor \frac{m}{2} + 1 \rfloor} d_{m,j}(a) \lambda_n^{\frac{1}{2} + \frac{1-j}{m}} + o(1) = \frac{(2n + 1) \pi}{2K_{m,0} \sin\left(\frac{2\pi}{m}\right)}.$$

Introduce the decomposition $\lambda_n = \lambda_{0,n} + \lambda_{1,n}$, where

$$\lambda_{0,n} = \left(\frac{(2n+1)\pi}{2K_{m,0} \sin\left(\frac{2\pi}{m}\right)} \right)^{\frac{2m}{m+2}} \quad \text{and} \quad \frac{\lambda_{1,n}}{\lambda_{0,n}} = o(1).$$

Then from (37) we have

$$\begin{aligned} \lambda_{0,n}^{\frac{1}{2}+\frac{1}{m}} &= \lambda_{0,n}^{\frac{1}{2}+\frac{1}{m}} \left(1 + \frac{\lambda_{1,n}}{\lambda_{0,n}} \right)^{\frac{1}{2}+\frac{1}{m}} + \sum_{j=1}^{\lfloor \frac{m}{2}+1 \rfloor} d_{m,j}(a) \lambda_{0,n}^{\frac{1}{2}+\frac{1-j}{m}} \left(1 + \frac{\lambda_{1,n}}{\lambda_{0,n}} \right)^{\frac{1}{2}+\frac{1-j}{m}} + o(1) \\ &= \lambda_{0,n}^{\frac{1}{2}+\frac{1}{m}} \left(1 + \sum_{k=1}^{\infty} \binom{\frac{1}{2}+\frac{1}{m}}{k} \left(\frac{\lambda_{1,n}}{\lambda_{0,n}} \right)^k \right) \\ &\quad + \sum_{j=1}^{\lfloor \frac{m}{2}+1 \rfloor} d_{m,j}(a) \lambda_{0,n}^{\frac{1}{2}+\frac{1-j}{m}} \left(1 + \sum_{k=1}^{\infty} \binom{\frac{1}{2}+\frac{1-j}{m}}{k} \left(\frac{\lambda_{1,n}}{\lambda_{0,n}} \right)^k \right) + o(1). \end{aligned}$$

Thus,

$$\begin{aligned} 0 &= \left(\frac{1}{2} + \frac{1}{m} \right) \frac{\lambda_{1,n}}{\lambda_{0,n}} + \sum_{k=2}^{\infty} \binom{\frac{1}{2}+\frac{1}{m}}{k} \left(\frac{\lambda_{1,n}}{\lambda_{0,n}} \right)^k \\ &\quad + \sum_{j=1}^{\lfloor \frac{m}{2}+1 \rfloor} d_{m,j}(a) \lambda_{0,n}^{-\frac{j}{m}} \left(1 + \sum_{k=1}^{\infty} \binom{\frac{1}{2}+\frac{1-j}{m}}{k} \left(\frac{\lambda_{1,n}}{\lambda_{0,n}} \right)^k \right) + o\left(\lambda_{0,n}^{-\frac{1}{2}-\frac{1}{m}} \right), \end{aligned}$$

and hence

$$\begin{aligned} &\frac{\lambda_{1,n}}{\lambda_{0,n}} + \frac{2m}{m+2} \sum_{k=2}^{\infty} \binom{\frac{1}{2}+\frac{1}{m}}{k} \left(\frac{\lambda_{1,n}}{\lambda_{0,n}} \right)^k \\ &\quad + \frac{2m}{m+2} \sum_{j=1}^{\lfloor \frac{m}{2}+1 \rfloor} d_{m,j}(a) \lambda_{0,n}^{-\frac{j}{m}} \left(\sum_{k=1}^{\infty} \binom{\frac{1}{2}+\frac{1-j}{m}}{k} \left(\frac{\lambda_{1,n}}{\lambda_{0,n}} \right)^k \right) + o\left(\lambda_{0,n}^{-\frac{1}{2}-\frac{1}{m}} \right) \\ (38) \quad &= -\frac{2m}{m+2} \sum_{j=1}^{\lfloor \frac{m}{2}+1 \rfloor} d_{m,j}(a) \lambda_{0,n}^{-\frac{j}{m}}. \end{aligned}$$

Thus, one concludes $\frac{\lambda_{1,n}}{\lambda_{0,n}} = \lambda_{2,n} + \lambda_{3,n}$, where

$$(39) \quad \lambda_{2,n} = -\frac{2m}{m+2} d_{m,1}(a) \lambda_{0,n}^{-\frac{1}{m}} \quad \text{and} \quad \lambda_{3,n} = o\left(\lambda_{0,n}^{-\frac{1}{m}} \right).$$

Next, from (39) along with (38) we have

$$\begin{aligned} &\lambda_{2,n} + \lambda_{3,n} + \frac{2m}{m+2} \sum_{k=2}^{\infty} \binom{\frac{1}{2}+\frac{1}{m}}{k} (\lambda_{2,n} + \lambda_{3,n})^k \\ &\quad + \frac{2m}{m+2} \sum_{j=1}^{\lfloor \frac{m}{2}+1 \rfloor} d_{m,j}(a) \lambda_{0,n}^{-\frac{j}{m}} \left(\sum_{k=1}^{\infty} \binom{\frac{1}{2}+\frac{1-j}{m}}{k} (\lambda_{2,n} + \lambda_{3,n})^k \right) + o\left(\lambda_{0,n}^{-\frac{1}{2}-\frac{1}{m}} \right) \\ (40) \quad &= -\frac{2m}{m+2} \sum_{j=1}^{\lfloor \frac{m}{2}+1 \rfloor} d_{m,j}(a) \lambda_{0,n}^{-\frac{j}{m}}. \end{aligned}$$

Thus,

$$\begin{aligned}
& \lambda_{3,n} + \frac{2m}{m+2} \sum_{k=2}^{\infty} \binom{\frac{1}{2} + \frac{1}{m}}{k} \sum_{\ell=0}^k \binom{k}{\ell} \lambda_{2,n}^{\ell} \lambda_{3,n}^{k-\ell} \\
& + \frac{2m}{m+2} \sum_{j=1}^{\lfloor \frac{m}{2} + 1 \rfloor} d_{m,j}(a) \lambda_{0,n}^{-\frac{j}{m}} \left(\sum_{k=1}^{\infty} \binom{\frac{1}{2} + \frac{1-j}{m}}{k} \sum_{\ell=0}^k \binom{k}{\ell} \lambda_{2,n}^{\ell} \lambda_{3,n}^{k-\ell} \right) + o\left(\lambda_{0,n}^{-\frac{1}{2} - \frac{1}{m}}\right) \\
(41) \quad & = -\frac{2m}{m+2} \sum_{j=2}^{\lfloor \frac{m}{2} + 1 \rfloor} d_{m,j}(a) \lambda_{0,n}^{-\frac{j}{m}},
\end{aligned}$$

and hence

$$\begin{aligned}
& \lambda_{3,n} + \frac{2m}{m+2} \sum_{k=2}^{\infty} \binom{\frac{1}{2} + \frac{1}{m}}{k} \sum_{\ell=0}^{k-1} \binom{k}{\ell} \lambda_{2,n}^{\ell} \lambda_{3,n}^{k-\ell} \\
& + \frac{2m}{m+2} \sum_{j=1}^{\lfloor \frac{m}{2} + 1 \rfloor} d_{m,j}(a) \lambda_{0,n}^{-\frac{j}{m}} \left(\sum_{k=1}^{\infty} \binom{\frac{1}{2} + \frac{1-j}{m}}{k} \sum_{\ell=0}^{k-1} \binom{k}{\ell} \lambda_{2,n}^{\ell} \lambda_{3,n}^{k-\ell} \right) + o\left(\lambda_{0,n}^{-\frac{1}{2} - \frac{1}{m}}\right) \\
& = -\frac{2m}{m+2} \sum_{j=2}^{\lfloor \frac{m}{2} + 1 \rfloor} d_{m,j}(a) \lambda_{0,n}^{-\frac{j}{m}} - \frac{2m}{m+2} \sum_{k=2}^{\infty} \binom{\frac{1}{2} + \frac{1}{m}}{k} \lambda_{2,n}^k \\
(42) \quad & - \frac{2m}{m+2} \sum_{j=1}^{\lfloor \frac{m}{2} + 1 \rfloor} d_{m,j}(a) \lambda_{0,n}^{-\frac{j}{m}} \left(\sum_{k=1}^{\infty} \binom{\frac{1}{2} + \frac{1-j}{m}}{k} \lambda_{2,n}^k \right).
\end{aligned}$$

So we choose

$$(43) \quad \lambda_{3,n} = \lambda_{4,n} + \lambda_{5,n},$$

where

$$\begin{aligned}
\lambda_{4,n} &= -\frac{2m}{m+2} d_{m,2}(a) \lambda_{0,n}^{-\frac{2}{m}} - \frac{2m}{m+2} \binom{\frac{1}{2} + \frac{1}{m}}{2} \lambda_{2,n}^2 - \frac{m d_{m,1}(a)}{m+2} \lambda_{0,n}^{-\frac{1}{m}} \lambda_{2,n} \\
&= \left(-\frac{2m}{m+2} d_{m,2}(a) + \left(\frac{2m^2}{(m+2)^2} - \left(\frac{2m}{m+2} \right)^3 \binom{\frac{1}{2} + \frac{1}{m}}{2} \right) d_{m,1}(a)^2 \right) \lambda_{0,n}^{-\frac{2}{m}}, \\
\lambda_{5,n} &= o\left(\lambda_{0,n}^{-\frac{2}{m}}\right).
\end{aligned}$$

Next, we replace $\lambda_{3,n}$ in (42) by (43). Upon iterating this process we get

$$\begin{aligned}
\lambda_n &= \lambda_{0,n} + \lambda_{1,n} = \lambda_{0,n} \left(1 + \frac{\lambda_{1,n}}{\lambda_{0,n}} \right) \\
&= \lambda_{0,n} (1 + \lambda_{2,n} + \lambda_{3,n}) \\
&= \lambda_{0,n} (1 + \lambda_{2,n} + \lambda_{4,n} + \lambda_{5,n}) \\
&\dots \\
(44) \quad &= \lambda_{0,n} \left(1 + \sum_{\ell=1}^{\lfloor \frac{m}{2} + 1 \rfloor} e_{\ell}(a) \lambda_{0,n}^{-\frac{\ell}{m}} + o\left(\lambda_{0,n}^{-\frac{1}{2} - \frac{1}{m}}\right) \right),
\end{aligned}$$

as $n \rightarrow +\infty$, that is, (5).

Suppose that $m = 3$. For this case we will use the asymptotic expansion in Theorem 14 that is valid in (27). Similarly to what we did for the case $m \geq 4$, if $C(a, \lambda) = 0$ then from the asymptotic expansion in Theorem 14 we have

$$[1 + o(1)] \exp [L(G^4(a), \omega^{-2}\lambda) + L(G^2(a), \omega^{-1}\lambda)] = -i\omega^{\frac{15}{4}}.$$

Thus, since $L(a, \lambda) = K_{3,0}(a)\lambda^{\frac{5}{6}} + K_{3,1}(a)\lambda^{\frac{3}{6}} + K_{3,2}(a)\lambda^{\frac{1}{6}} + o(1)$, we have

$$\begin{aligned} & L(G^4(a), \omega^{-2}\lambda) + L(G^2(a), \omega^{-1}\lambda) + o(1) \\ &= K_{3,0}(G^4(a)) (\omega^{-2}\lambda)^{\frac{5}{6}} + K_{3,1}(G^4(a)) (\omega^{-2}\lambda)^{\frac{3}{6}} + K_{3,2}(G^4(a)) (\omega^{-2}\lambda)^{\frac{1}{6}} \\ &+ K_{3,0}(G^2(a)) (\omega^{-1}\lambda)^{\frac{5}{6}} + K_{3,1}(G^2(a)) (\omega^{-1}\lambda)^{\frac{3}{6}} + K_{3,2}(G^2(a)) (\omega^{-1}\lambda)^{\frac{1}{6}} + o(1) \\ &= K_{3,0} \left(e^{-i\frac{2\pi}{3}} + e^{-i\frac{\pi}{3}} \right) \lambda^{\frac{5}{6}} + c_{3,1}(a)\lambda^{\frac{3}{6}} + c_{3,2}(a)\lambda^{\frac{1}{6}} + o(1) \\ &= -2iK_{3,0} \sin \left(\frac{2\pi}{3} \right) \lambda^{\frac{5}{6}} + c_{3,1}(a)\lambda^{\frac{3}{6}} + c_{3,2}(a)\lambda^{\frac{1}{6}} + o(1). \end{aligned}$$

So the continuous function $\lambda \mapsto L(G^4(a), \omega^{-2}\lambda) + L(G^2(a), \omega^{-1}\lambda) + o(1)$ maps a neighborhood of the positive real axis near infinity onto a neighborhood of the negative imaginary axis near infinity. Hence, there exist a sequence of λ_n near the positive real axis such that for all large enough positive integers n ,

$$-2iK_{3,0} \sin \left(\frac{2\pi}{3} \right) \lambda_n^{\frac{5}{6}} + c_{3,1}(a)\lambda_n^{\frac{3}{6}} + c_{3,2}(a)\lambda_n^{\frac{1}{6}} + o(1) = \ln \left(-i\omega^{\frac{15}{4}} \right) = (\pi - 2(n+1)\pi) i.$$

From this result one concludes that the asymptotic expansion (5) holds for $m = 3$ as well similarly to the proof for the case $m \geq 4$. $\square$

6. PROOF OF THEOREM 3

Proof of Theorem 3. First, note from (5) that $\arg(\lambda_n) \rightarrow 0$ as $n \rightarrow +\infty$.

Next, we have

$$\begin{aligned} \lambda_{0,n+1} &= \left(\frac{(2n+3)\pi}{2K_{m,0} \sin \left(\frac{2\pi}{m} \right)} \right)^{\frac{2m}{m+2}} \\ &= \left(\frac{(2n+1)\pi}{2K_{m,0} \sin \left(\frac{2\pi}{m} \right)} + \frac{2\pi}{2K_{m,0} \sin \left(\frac{2\pi}{m} \right)} \right)^{\frac{2m}{m+2}} \\ &= \lambda_{0,n} \left(1 + \frac{2}{2n+1} \right)^{\frac{2m}{m+2}} \\ &= \lambda_{0,n} \left(1 + \frac{2m}{m+2} \frac{2}{2n+1} + O \left(\frac{1}{n^2} \right) \right) \\ (45) \quad &= \lambda_{0,n} + \frac{2m\pi}{(m+2)K_{m,0} \sin \left(\frac{2\pi}{m} \right)} \lambda_{0,n}^{1-\frac{1}{2}-\frac{1}{m}} + o \left(\lambda_{0,n}^{\frac{1}{2}-\frac{1}{m}} \right). \end{aligned}$$

Thus,

$$\lambda_{n+1} - \lambda_n \underset{n \rightarrow +\infty}{=} \frac{2m\pi}{(m+2)K_{m,0} \sin\left(\frac{2\pi}{m}\right)} \lambda_{0,n}^{\frac{1}{2}-\frac{1}{m}} + o\left(\lambda_{0,n}^{\frac{1}{2}-\frac{1}{m}}\right),$$

and hence, $|\lambda_{n+1} - \lambda_n| \rightarrow \infty$ and $\arg(\lambda_{n+1} - \lambda_n) \rightarrow 0$ as $n \rightarrow +\infty$. Since $\arg(\lambda_n) \rightarrow 0$ (and $\arg(\lambda_{n+1}) \rightarrow 0$) as $n \rightarrow +\infty$, there exists $N \in \mathbb{N}$ such that $|\lambda_n| < |\lambda_{n+1}|$ if $n \geq N$. $\square$

Remark. Here we will show that if $a \in \mathbb{R}^{m-1}$, then $e_\ell(a) \in \mathbb{R}$ for all $1 \leq \ell \leq \frac{m}{2} + 1$ with $e_\ell(a)$ defined in (44).

From (14) one can see that $\overline{K_{m,j}(G^{-1}(\bar{a}))} = K_{m,j}(G(a))$. Next, suppose that $a \in \mathbb{R}^{m-1}$. If $m \geq 4$ then from (34),

$$\begin{aligned} ic_{m,j}(a) &= i \left(K_{m,j}(G(a))(\omega^2)^{\frac{1}{2}+\frac{1-j}{m}} - K_{m,j}(G^{-1}(a))(\omega^{-2})^{\frac{1}{2}+\frac{1-j}{m}} \right) \\ &= i \left(K_{m,j}(G(a))(\omega^2)^{\frac{1}{2}+\frac{1-j}{m}} - \overline{K_{m,j}(G(a))(\omega^2)^{\frac{1}{2}+\frac{1-j}{m}}} \right) \in \mathbb{R}, \quad 1 \leq j \leq \frac{m}{2} + 1. \end{aligned}$$

So by (36), $d_{m,j}(a) \in \mathbb{R}$ for all $1 \leq j \leq \frac{m}{2} + 1$, and hence by (44), $e_\ell(a) \in \mathbb{R}$ for all $1 \leq \ell \leq \frac{m}{2} + 1$.

If $m = 3$ then one can show $e_\ell(a) \in \mathbb{R}$ for $\ell = 1, 2$, using the formulas at the end of the proof of Theorem 2.

Acknowledgments. The author thanks Fritz Gesztesy and Richard Laugesen for critical reading of this manuscript and suggestions.

REFERENCES

- [1] C. M. Bender, M. Berry, P. N. Meisinger, V. M. Savage and M. Simsek Complex WKB Analysis of Energy-Level Degeneracies of Non-Hermitian Hamiltonians, *J. Phys. A: Math. Gen.*, 34:L31–L36, 2001.
- [2] C. M. Bender and S. Boettcher. Real spectra in non-Hermitian Hamiltonians having $\mathcal{PT}$ -symmetry. *Phys. Rev. Lett.*, 80:5243–5246, 1998.
- [3] C. M. Bender and A. Turbinger. Analytic continuation of eigenvalue problems. *Phys. Lett. A*, 173:442–446, 1993.
- [4] C. M. Bender and E. J. Weniger. Numerical evidence that the perturbation expansion for a non-Hermitian $\mathcal{PT}$ -symmetric Hamiltonian is Stieltjes. *J. Math. Phys.*, 42:2167–2183, 2001.
- [5] E. Caliceti, S. Graffi and M. Maioli. Perturbation theory of odd anharmonic oscillators. *Comm. Math. Phys.*, 75:51–66, 1980.
- [6] E. Delabaere and F. Pham. Eigenvalues of complex Hamiltonians with $\mathcal{PT}$ -symmetry I, II. *Phys. Lett. A*, 250:25–32, 1998.
- [7] E. Delabaere and D. T. Trinh. Spectral analysis of the complex cubic oscillator. *J. Phys. A: Math. Gen.*, 33:8771–8796, 2000.
- [8] P. Dorey, C. Dunning and R. Tateo. Spectral equivalences, Bethe ansatz equations, and reality properties in $\mathcal{PT}$ -symmetric quantum mechanics. *J. Phys. A: Math. Gen.*, 34:5679–5704, 2001.
- [9] P. Dorey and R. Tateo. On the relation between Stokes multipliers and T - Q systems of conformal field theory. *Nucl. Phys. B*, 563:573–602, 1999.
- [10] C. R. Handy. Generating converging bounds to the (complex) discrete states of the $P^2 + iX^3 + i\alpha X$ Hamiltonian. *J. Phys. A: Math. Gen.*, 34:5065–5081, 2001.
- [11] C. R. Handy, D. Khan, Xiao-Qian Wang and C. J. Tymczak. Multiscale reference function analysis of the $\mathcal{PT}$ symmetry breaking solutions for the $P^2 + iX^3 + i\alpha X$ Hamiltonian. *J. Phys. A: Math. Gen.*, 34:5593–5602, 2001.
- [12] E. Hille. *Lectures on Ordinary Differential Equations*. Addison-Wesley, Reading, Massachusetts, 1969.

- [13] G. A. Mezincescu. Some properties of eigenvalues and eigenfunctions of the cubic oscillator with imaginary coupling constant. *J. Phys. A: Math. Gen.*, 33:4911–4916, 2000.
- [14] A. Mostafazadeh. Pseudo-Hermiticity versus PT Symmetry: The necessary condition for the reality of the spectrum of a non-Hermitian Hamiltonian. *J. Math. Phys.*, 43:205–214, 2002.
- [15] K. C. Shin. On the eigenproblems of $\mathcal{PT}$ -symmetric oscillators. *J. Math. Phys.*, 42:2513–2530, 2001.
- [16] K. C. Shin. On the reality of the eigenvalues for a class of $\mathcal{PT}$ -symmetric oscillators, *Comm. Math. Phys.*, 229(3):543–564, 2002.
- [17] B. Simon. Coupling Constant Analyticity for the Anharmonic Oscillator. *Ann. Phys.*, 58:76–136, 1970.
- [18] Y. Sibuya. *Global theory of a second order linear ordinary differential equation with a polynomial coefficient*. North-Holland Publishing Company, Amsterdam-Oxford, 1975.
- [19] M. Znojil. Spiked and $\mathcal{PT}$ -symmetrized decadic potentials supporting elementary N -plets of bound states. *J. Phys. A: Math. Gen.*, 33:4911–4916, 2000.

EMAIL CONTACT: kcshin@math.missouri.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211, USA